

Mass-number and isotope dependence of the local microscopic optical potential for polarized proton scattering

Masakazu Toyokawa,^{1,*} Kosho Minomo,^{1,†} and Masanobu Yahiro^{1,‡}

¹*Department of Physics, Kyushu University, Fukuoka 812-8581, Japan*

(Dated: May 1, 2013)

We derive the local microscopic optical potential U systematically for polarized proton scattering at 65 MeV, using the local-potential version of the Melbourne g -matrix folding model. As target nuclei, we take ${}^6\text{He}$ and neutron-rich Ne isotopes in addition to stable nuclei of mass number $A = 4\text{--}208$ in order to clarify A and isotope dependence of U . The local potential reproduces the experimental data systematically and has a geometry similar to the phenomenological optical potentials for stable targets. The target density is broadened by the weak-binding nature and/or the deformation in unstable nuclei. For the real part of the spin-orbit potential, the density broadening weakens the strength and enlarges the radius, whereas for the central potential it enlarges both the strength and the radius. The density-broadening effect is conspicuous for halo nuclei such as ${}^6\text{He}$ and ${}^{31}\text{Ne}$. Similar discussion is made briefly for proton scattering at 200 MeV.

PACS numbers: 25.40.Cm, 24.10.Ht, 24.70.+s

I. INTRODUCTION

Understanding of nucleon-nucleus (NA) and nucleus-nucleus (AA) interactions are one of the most important subjects in nuclear physics. The interactions (optical potentials) are necessary to describe not only elastic scattering but also more complicated reactions. Actually the interactions are inputs of theoretical calculations such as the distorted-wave Born approximation (DWBA) and the continuum discretized coupled-channels method (CDCC) [1–3] to analyze inelastic scattering and transfer and breakup reactions.

In general, NA elastic scattering is less absorptive and hence more sensitive to nuclear interior than AA scattering. Furthermore, one can determine not only the central part but also the spin-orbit part of NA interaction, when the incident particle is polarized. From this point of view, systematic measurements of polarized proton (p) elastic scattering were made so far for stable target nuclei. As an outcome of the measurements, some global NA optical potentials have been constructed phenomenologically; see for example Refs. [4–7]. In most of the cases, the potentials are assumed to be local and thereby quite practical in many applications such as DWBA and CDCC calculations.

Elastic scattering of unstable nuclei on a polarized proton target is a quite good tool of investigating nuclear structures of unstable nuclei and interactions between a proton and unstable nuclei, but the measurement is not easy. The first measurement on the vector analyzing power A_y was made by Uesaka and his collaborators for $p+{}^6\text{He}$ elastic scattering at 71 MeV with the inverse kinematics [8]. The g -matrix folding model yields a rea-

sonable agreement with the measured vector analyzing power [8]. The nuclear-medium effect is thus important for the observable. Very lately the effect of ${}^6\text{He}$ breakup on the elastic scattering was investigated with the eikonal approximation [9]. The medium effect is more significant than the breakup effect for A_y .

The g -matrix folding model is a powerful tool to describe NA and AA interactions [10–19]. Particularly when the Melbourne g -matrix nucleon-nucleon (NN) interaction [17] is used, the model is successful in reproducing polarized proton elastic scattering from stable target nuclei systematically with no free parameter [17]. In the model, effects of target excitation and breakup in addition to the medium effect are taken into account within the local-density approximation. The microscopic optical potential derived by the model is nonlocal and thereby not so practical in many applications. It is, however, possible to localize the potential with the Brieda-Rook approximation [12]. Recently the validity of the approximation was shown in Ref. [20].

From a theoretical viewpoint based on the multiple scattering theory [21–23], a multiple NN scattering series in AA collision [23] is more complicated than that in NA collision [21, 22]. In this sense, microscopic understanding of AA interaction is relatively more difficult than that of NA interaction. Very recently, however, for AA elastic scattering the local microscopic optical potential was derived from the Melbourne g -matrix interaction by using both the folding model and the Brieda-Rook localization [24–26]. When the local-potential version of the Melbourne g -matrix folding model was applied to the ${}^{12}\text{C}+{}^{12}\text{C}$ scattering at 6.2, 135 and 250 MeV/nucleon and ${}^{20\text{--}32}\text{Ne}+{}^{12}\text{C}$ scattering at 240 MeV/nucleon, the resultant local potentials reproduced the measured elastic cross sections and reaction cross sections σ_R with no free parameter [24–26]. In the calculation, the densities of unstable nuclei ${}^{20\text{--}32}\text{Ne}$ were evaluated by antisymmetrized molecular dynamics (AMD) [27, 28] with the Gogny-D1S interaction [29], whereas the phenomenolog-

*toyokawa@phys.kyushu-u.ac.jp

†minomo@phys.kyushu-u.ac.jp

‡yahiro@phys.kyushu-u.ac.jp

ical density [30] was taken for ^{12}C . The AMD wave functions successfully describes low-lying spectra of Ne isotopes [27]. The microscopic approach shows that $^{30-32}\text{Ne}$ in the “island of inversion” have large deformation and particularly ^{31}Ne is a deformed halo nucleus with spin-parity $3/2^-$ [24–26]. The local-potential version of the Melbourne g -matrix folding model is thus a powerful tool of determining the structure of unstable nuclei. The success of the local-potential version of the Melbourne g -matrix folding model for proton scattering from stable nuclei and $^{20-32}\text{Ne}+^{12}\text{C}$ scattering implies that the model is reliable also for proton scattering from unstable nuclei such as Ne isotopes.

In this paper, we derive the local microscopic optical potential U systematically for polarized proton scattering at 65 MeV, using the local-potential version of the Melbourne g -matrix folding model. This local nature of U makes it easy to clarify global properties of U for both stable and unstable targets. As the targets, we consider ^6He and neutron-rich Ne isotopes in addition to stable nuclei of mass number $A = 4-208$. Throughout the global analyses, we clarify A and isotope dependence of U . The target density is broadened much in ^6He and neutron-rich Ne isotopes compared with stable nuclei with the same A by the weak-binding nature and/or the nuclear deformation. We clarify how the density broadening affects the shape and the strength of U . This analysis is interesting particularly for ^6He and ^{31}Ne , since ^6He is a typical two-neutron halo nucleus and ^{31}Ne is a one-neutron halo nucleus with large deformation. Similar analyses are briefly made for proton scattering at 200 MeV.

We recapitulate the local-potential version of the Melbourne g -matrix folding model in Sec. II and show numerical results in Sec. III. Section IV is devoted to summary.

II. THEORETICAL FRAMEWORK

Proton elastic scattering can be described by the one-body Schrödinger equation

$$(T_R + U - E)\Psi^{(+)} = 0, \quad (1)$$

with the proton optical potential U , where E is the incident energy of proton and T_R is the kinetic energy operator concerning the relative coordinate $\mathbf{R}$ between an incident proton and a target. The optical potential U can be divided into the central (CE), the spin-orbit (LS) and the Coulomb (Coul) component:

$$U = U_{\text{CE}} + U_{\text{LS}}\mathbf{L} \cdot \boldsymbol{\sigma} + V_{\text{Coul}}. \quad (2)$$

In the g -matrix folding model, U consists of the direct and exchange parts [31]. The exchange part is nonlocal, but it can be localized with the Brieva-Rook approximation [12]. The validity of the approximation is shown in Ref. [20]. We take this approximation in the present paper, since the local nature of U makes it possible to make global analyses of U over stable and unstable targets. In

addition to these merits, the local microscopic optical potential is quite useful in many applications, particularly when a potential between stable and unstable nuclei is needed.

The central part U_{CE} is then described by the sum of the direct component $U_{\text{CE}}^{\text{DR}}$ and the localized exchange component $U_{\text{CE}}^{\text{EX}}$ [12, 15, 18]:

$$U_{\text{CE}} \equiv V_{\text{CE}} + iW_{\text{CE}} = U_{\text{CE}}^{\text{DR}} + U_{\text{CE}}^{\text{EX}} \quad (3)$$

with

$$U_{\text{CE}}^{\text{DR}}(\mathbf{R}) = \sum_{\alpha=p,n} \int \rho_{\alpha}(\mathbf{r}) g_{p\alpha}^{\text{DR}}(s; \rho_{\alpha}) d\mathbf{r}, \quad (4)$$

$$U_{\text{CE}}^{\text{EX}}(\mathbf{R}) = - \sum_{\alpha=p,n} \int \rho_{\alpha}(\mathbf{r}, \mathbf{r} - \mathbf{s}) g_{p\alpha}^{\text{EX}}(s; \rho_{\alpha}) j_0(K(R)s) d\mathbf{r}, \quad (5)$$

where V_{CE} (W_{CE}) is the real (imaginary) part of U_{CE} and $\mathbf{s} = \mathbf{r} - \mathbf{R}$ for coordinate $\mathbf{r}$ of an interacting nucleon from the center-of-mass (c.m.) of target. The local momentum $\hbar K(R) \equiv \sqrt{2\mu_R(E - U_{\text{CE}} - V_{\text{Coul}})}$ present in Eq. (5) is obtained self-consistently, where μ_R is the reduced mass of the proton-target system.

Usually the direct part $g_{p\alpha}^{\text{DR}}$ and the exchange part $g_{p\alpha}^{\text{EX}}$ of the g -matrix interaction are assumed to be a function of the local density $\rho_{\alpha} = \rho_{\alpha}(\mathbf{r} - \mathbf{s}/2)$ at the midpoint of the interacting nucleon pair. The direct and exchange parts are described as

$$g_{pp}^{\text{DR,EX}}(s; \rho_p) = \frac{1}{4} (\pm g^{01} + 3g^{11}), \quad (6)$$

$$g_{pn}^{\text{DR,EX}}(s; \rho_n) = \frac{1}{8} (g^{00} \pm g^{01} \pm 3g^{10} + 3g^{11}), \quad (7)$$

in terms of the spin-isospin component g^{ST} of the g -matrix interaction.

Similar derivation is possible for the spin-orbit part U_{LS} [18]:

$$U_{\text{LS}} \equiv V_{\text{LS}} + iW_{\text{LS}} = U_{\text{LS}}^{\text{DR}} + U_{\text{LS}}^{\text{EX}} \quad (8)$$

with

$$U_{\text{LS}}^{\text{DR}}(\mathbf{R}) = -\frac{1}{4R^2} \sum_{\alpha=p,n} \int \mathbf{R} \cdot \mathbf{s} \rho_{\alpha}(\mathbf{r}) g_{\text{LS},p\alpha}^{\text{DR}}(s; \rho_{\alpha}) d\mathbf{r}, \quad (9)$$

$$U_{\text{LS}}^{\text{EX}}(\mathbf{R}) = -\pi \sum_{\alpha=p,n} \int ds s^3 \left[\frac{2j_0(K(R)s)}{R} \rho_1^{(\alpha)}(R, s) + \frac{j_1(K(R)s)}{2k} \delta_0^{(\alpha)}(R, s) \right], \quad (10)$$

where V_{LS} (W_{LS}) is the real (imaginary) part of U_{LS} and see Appendix A for the definition of $\rho_1^{(\alpha)}(R, s)$ and $\delta_0^{(\alpha)}(R, s)$. As the g -matrix interaction, we take the Melbourne interaction [17] constructed from the Bonn-B nucleon-nucleon potential [32].

As the target density ρ , we consider three kinds of matter densities. For all the targets, the matter densities are evaluated by spherical Hartree-Fock (HF) calculations with the Gogny-D1S interaction [29] in which

the spurious c.m. motion is removed in the standard manner [26]. For lighter stable nuclei of $A \leq 40$, we also use the phenomenological proton-density [30] determined from the electron scattering in which the finite-size effect of the proton charge is unfolded in the usual way [33]. The neutron density is assumed to have the same geometry as the corresponding proton one, since the deviation of the neutron root-mean-square (RMS) radius from the proton one is about 1% in the spherical HF calculation.

For $^{20-32}\text{Ne}$, the matter densities are evaluated also by AMD [27, 28] with the Gogny-D1S interaction [29]. This provides the deformed matter densities. The AMD wave functions successfully describes low-lying spectra of Ne isotopes [27] and σ_R of $^{20-32}\text{Ne} + ^{12}\text{C}$ scattering at 240 MeV/nucleon [24–26], and it is free from the spurious c.m. motion; see Ref. [26, 27] for the details of AMD calculation.

Particularly for ^{31}Ne with quite small one-nucleon separation energy, we have to do a sophisticated calculation to make a tail correction to the AMD density, since the AMD density is inaccurate in its tail region. In principle, the ground state of ^{31}Ne can be described by the $^{30}\text{Ne} + n$ cluster model with core (^{30}Ne) excitations, and one can solve coupled equations for the model with the resonating group method (RGM) in which the ground and excited states of ^{30}Ne were constructed by AMD. This was done in Ref. [25, 26]. This method is referred to as AMD-RGM in this paper.

The matter density $\rho_{ImIm'}(\mathbf{r})$ is obtainable from the ground state wave function $\Phi_{\text{g.s.}}^{Im\pi}$ with spin I , its z -component m and parity π as

$$\rho_{ImIm'}(\mathbf{r}) = \langle \Phi_{\text{g.s.}}^{Im\pi} | \sum_i \delta(\mathbf{r}_i - \mathbf{X} - \mathbf{r}) | \Phi_{\text{g.s.}}^{Im'\pi} \rangle, \quad (11)$$

$$= \sum_{\lambda=0}^{2I} \rho_{II}^{(\lambda)}(r) (Im' \lambda \mu | Im) Y_{\lambda\mu}^*(\hat{r}), \quad (12)$$

where $\mathbf{X}$ is the c.m. coordinate and $\mathbf{r}_i$ is the coordinate of i -th nucleon, and the summation of λ in Eq. (12) runs for even numbers. As a standard manner in the folding model, only $\rho_{II}^{(\lambda=0)}$ is taken. This procedure is exact for even nuclei with $I = 0$, but approximate for odd nuclei with $I \neq 0$. It is, however, confirmed in Ref. [26] that the procedure is a quite good approximation.

III. RESULTS

In this section we analyze polarized proton scattering at $E = 65$ and 200 MeV, using the local-potential version of the Melbourne g -matrix folding model, although the analysis is mainly focused on the case of $E = 65$ MeV. In the folding model, we also consider the Franey-Love t -matrix NN interaction [34]. As target nuclei of proton scattering, we consider stable nuclei of ^4He , ^{12}C , ^{16}O , ^{20}Ne , ^{24}Mg , ^{40}Ca , ^{90}Zr , ^{208}Pb and unstable nuclei of ^6He and neutron-rich Ne isotopes.

A. Proton scattering from stable nuclei at 65 and 200 MeV

In this subsection, we mainly consider proton scattering at 65 MeV and discuss the case of 200 MeV at the end of this subsection.

Figure 1 shows the elastic cross section $d\sigma/d\Omega$ and the vector analyzing power A_y as a function of the scattering angle $\theta_{\text{c.m.}}$ for proton scattering at $E = 65$ MeV from ^4He , ^{12}C , ^{40}Ca and ^{208}Pb targets. The solid lines denote results of the local microscopic potential derived from the Melbourne g -matrix interaction. The results are consistent with those [17] of the nonlocal microscopic potential derived from the Melbourne g -matrix interaction, although the target densities used are different between the two calculations. The g -matrix folding model (solid line) yields better agreement with the experimental data [35–42] than the t -matrix folding model (dashed line). The medium effect shown by the difference between the solid and dashed lines are more significant for A_y . For lighter targets of ^4He , ^{12}C and ^{40}Ca , the agreement of the g -matrix folding model with the data is better for the phenomenological density (solid line) than for the HF density (dotted line). However, the g -matrix folding model with the HF density still keeps reasonable agreement with the data. The HF density based on the Gogny-D1S interaction [29] is thus quite useful to derive the microscopic optical potential U systematically.

Figure 2 shows reaction cross sections σ_R as a function of A . Closed circles (cross symbols) denote results of the g -matrix (t -matrix) folding model with the phenomenological target density for $A \leq 40$ and the HF target density for $A > 40$. The g -matrix folding model yields better agreement with the data than the t -matrix folding model. The results of the g -matrix folding model agree with the data [43–47] within 10% error in a wide range of $4 \leq A \leq 208$. Since σ_R is sensitive to W_{CE} , the part is thus derived with at least 90% accuracy.

As a merit of deriving the local microscopic optical potential U , we can directly compare U with the phenomenological optical potentials. Here we take two types of the phenomenological optical potentials. One is the Koning-Delaroche (KD) optical potential [4] based on the standard potential search in the framework of the Schrödinger equation, and another is the Dirac phenomenology optical potential [5, 6] based on the potential search in the framework of the Dirac equation. As an example, we compare the three optical potentials for the $p + ^{40}\text{Ca}$ system at $E = 65$ MeV in Fig. 3. The microscopic optical potential is calculated from the phenomenological target density. The imaginary spin-orbit part W_{LS} is about ten times as small as V_{LS} , so that it hardly affects the present elastic scattering. We then compare the three optical potentials for V_{CE} , W_{CE} and V_{LS} . The KD optical potential (dashed line) has similar R dependence to the Dirac phenomenology optical potential (dotted line). In the peripheral region, furthermore, R dependence of the two phenomenological optical




FIG. 1: (Color online) Angular distributions of the elastic cross section $d\sigma/d\Omega$ (a) and the vector analyzing power A_y (b) for proton scattering at $E = 65$ MeV from ^4He , ^{12}C , ^{40}Ca and ^{208}Pb targets. The solid (dotted) lines represent the results of the g -matrix folding model with the phenomenological (HF) density, whereas the dashed lines correspond to results of the t -matrix folding model with the phenomenological density. Experimental data are taken from Refs. [35–42].

potentials is similar to that of the microscopic one (solid line). At small R , the deviation of the microscopic optical potential from the phenomenological ones is small for V_{CE} , but becomes sizable for V_{LS} and W_{CE} . For V_{LS} the deviation little affects the measured $d\sigma/d\Omega$ and A_y up to 90° , but for W_{CE} the deviation slightly enhances σ_R and consequently overestimates the measured σ_R by




FIG. 2: (Color online) A dependence of reaction cross sections σ_R at $E \approx 65$ MeV. Closed circles (cross symbols) stand for results of the g -matrix (t -matrix) folding model with the phenomenological density for $A \leq 40$ and the HF one for $A > 40$. Experimental data are taken from Refs. [43–47], where $E = 47.9$ MeV for ^4He , 65.5 MeV for ^{12}C , ^{16}O , ^{40}Ca , ^{208}Pb , 47.0 MeV for ^{20}Ne , 48.0 MeV for ^{24}Mg and 60.8 MeV for ^{90}Zr .

several percent.




FIG. 3: (Color online) R dependence of the microscopic optical potential (solid line), the Koning-Delaroche (KD) optical potential (dashed line) [4] and the Dirac phenomenology optical potential (dotted line) [5, 6] for $p+^{40}\text{Ca}$ scattering at $E = 65$ MeV.

For the $p+^4\text{He}$ system, the g -matrix folding model does not reproduce the measured elastic cross section at backward angles $\theta_{c.m.} \gtrsim 90^\circ$. A possible origin of this deviation is the heavy-ion (^3H) exchange process that is not included in the folding model. The process generates a parity-dependent potential [48]. We then rewrite V_{CE} as

$$V_{CE} \rightarrow V_{CE}(1 + \alpha(-1)^L) \quad (13)$$

to incorporate the effect in the folding model phenomenologically. The parameter α is fitted to the experimental data [35, 36], and the resulting value is $\alpha = 0.13$. In Fig. 4, the solid and dashed lines show $d\sigma/d\Omega$ and A_y calculated by the g -matrix folding model with $\alpha = 0.13$ and 0, respectively. The effect of the parity-dependent potential is appreciable at intermediate angles $60^\circ \lesssim \theta_{c.m.} \lesssim 90^\circ$ and significant at backward angles at $\theta_{c.m.} \gtrsim 90^\circ$.




FIG. 4: (Color online) Effect of the parity-dependent potential on the elastic cross section $d\sigma/d\Omega$ (a) and the vector analyzing power A_y (b) for $p+^4\text{He}$ scattering at $E = 65$ MeV. The solid (dashed) line stands for the result of the g -matrix folding model with $\alpha = 0.13$ (0), where the phenomenological density is taken for ^4He . Experimental data are taken from Refs. [35, 36].

For the $p+^{12}\text{C}$ system, the measured A_y is not reproduced around $\theta_{c.m.} = 100^\circ$ by the g -matrix folding model. Even the phenomenological optical potentials [4–6] cannot reproduce it. The parity-dependent potential does not solve this problem, since the potential enhances the elastic cross section largely at $\theta_{c.m.} > 90^\circ$. Therefore this is an open problem. Except for the measured A_y around $\theta_{c.m.} = 100^\circ$ for the $p+^{12}\text{C}$ system, the g -matrix folding model yields good agreement with the data for targets of $A = 4$ –208.

Now we briefly analyze proton scattering at $E = 200$ MeV as an example of intermediate energies and a typical case of beam energies in Radioactive Isotope Beam Factory (RIBF).

Figure 5 shows angular distributions of $d\sigma/d\Omega$ and A_y for the scattering from stable nuclei ^4He , ^{12}C , ^{40}Ca and ^{208}Pb . The definition of lines is the same as in Fig. 1. The g -matrix folding model with the phenomenological density, shown by the solid line, reproduces the data [51–

54] particularly at forward angles $\theta_{c.m.} \lesssim 60^\circ$. ($p+^{12}\text{C}$ 65 MeV : $\theta = 128^\circ$, 200 MeV : $\theta = 60^\circ$) When Fig. 5 is compared with Fig. 1 as a function of the transferred momentum, $\theta_{c.m.} = 60^\circ$ in Fig. 5 corresponds to $\theta_{c.m.} = 128^\circ$ in Fig. 1. Therefore, $\theta_{c.m.} > 60^\circ$ in Fig. 5 correspond to very backward angles $\theta_{c.m.} > 128^\circ$ in Fig. 1. In this sense, it is natural that in Fig. 5 the solid lines are deviated from the data at $\theta_{c.m.} \gtrsim 60^\circ$ particularly for lighter targets of $A \leq 12$. Even at $\theta_{c.m.} \gtrsim 60^\circ$ for lighter targets of $A \leq 12$, the g -matrix folding model with the phenomenological density (solid line) yields better agreement with the data than that with the HF density (dotted line). The medium effect, shown by the difference between the solid and dashed lines, is still significant at $E = 200$ MeV.

B. $p+^6\text{He}$ scattering at 71 and 200 MeV

As a typical example of proton scattering from unstable targets, we first consider $p+^6\text{He}$ scattering at 71 MeV, since the experimental data is available not only for $d\sigma/d\Omega$ but also for A_y . The ^6He density is obtained by the $^4\text{He}+n+n$ orthogonality-condition model; see Refs. [49, 50] for the detail. As the ^4He -density in the model, we take either the phenomenological or the HF density. In Fig. 6, the solid (dashed) line shows $d\sigma/d\Omega$ and A_y calculated by the g -matrix (t -matrix) folding model with the phenomenological ^4He density, whereas the dotted line corresponds to the result of the g -matrix folding model with the HF ^4He density. The g -matrix folding model (solid line) yields better agreement with the data [8, 55] than the t -matrix folding model (dashed line). The solid line is consistent with a result of the nonlocal microscopic optical potential derived from the Melbourne g -matrix interaction [8], although the target densities used are different between the two calculations.

In Fig. 6, the difference between the dashed and dotted lines comes from that between the phenomenological and HF ^4He densities. Comparing Fig. 6 with Fig. 1, one can see that the difference of ^4He density yields smaller effect on $d\sigma/d\Omega$ and A_y in the $p+^6\text{He}$ scattering than in the $p+^4\text{He}$ scattering. The difference of ^4He density is thus partly masked by the density of extra two neutrons in the $p+^6\text{He}$ scattering.

In Fig. 6, the nuclear medium effect is shown by the difference between the solid and dashed lines, and the effect is such more significant for A_y . The difference mainly comes from the fact that W_{CE} calculated with the g -matrix folding model is much less absorptive than that with the t -matrix folding model; see Fig. 7 for R dependence of U .

As mentioned above, the g -matrix folding model yields better agreement with the data than the t -matrix folding model. The agreement is, however, not perfect particularly for A_y . The parity-dependent potential little works at forward angles $\theta < 90^\circ$ in which the experimental data is available. We then make the optical potential search




FIG. 5: (Color online) Angular distributions of the elastic cross section $d\sigma/d\Omega$ (a) and the vector analyzing power A_y (b) for proton scattering at $E = 200$ MeV from ^4He , ^{12}C , ^{40}Ca and ^{208}Pb targets. See Fig. 1 for the definition of lines. Experimental data are taken from Refs. [51–54].

to see what causes the difference between the theoretical and experimental results. Here we change the strength of U only as

$$U \rightarrow (f_V^{\text{CE}} V_{\text{CE}} + i f_W^{\text{CE}} W_{\text{CE}}) + (f_V^{\text{LS}} V_{\text{LS}} + i f_W^{\text{LS}} W_{\text{LS}}) \mathbf{L} \cdot \boldsymbol{\sigma} + V_{\text{Coul}} \quad (14)$$

with adjustable parameters $(f_V^{\text{CE}}, f_W^{\text{CE}}, f_V^{\text{LS}}, f_W^{\text{LS}})$. The best fit is obtained when $(f_V^{\text{CE}}, f_W^{\text{CE}}, f_V^{\text{LS}}, f_W^{\text{LS}}) = (0.7, 0.7, 0.9, 4.0)$, but $d\sigma/d\Omega$ and A_y at $\theta > 40^\circ$ are not




FIG. 6: (Color online) Angular distributions of the elastic cross section $d\sigma/d\Omega$ (a) and the vector analyzing power A_y (b) for $p+^6\text{He}$ scattering at $E = 71$ MeV. See Fig. 1 for the definition of lines. Experimental data are taken from Ref. [8, 55].

sensitive to changes of f_W^{CE} and f_V^{LS} and hence these parameters are not determined sharply.

In Fig. 8, the result (solid line) of the potential search is compared with the original result (dashed line). Small reduction of $d\sigma/d\Omega$ from the dashed to the solid line at $\theta > 50^\circ$ mainly comes from reduction of V_{CE} , and large change of A_y from the dashed to the solid line at $\theta > 40^\circ$ is mainly originated in large enhancement of W_{LS} . Thus the large enhancement of W_{LS} is necessary, but it is not clear what causes the large enhancement of W_{LS} . This is an interesting question to be solved in future.

Now we briefly analyze proton scattering from ^6He at $E = 200$ MeV. Experimental data is not available for this energy at the present stage, but the measurement of A_y is planned in RIBF. Figure 9 shows angular distributions of $d\sigma/d\Omega$ and A_y for proton scattering from ^6He at $E = 200$ MeV. The definition of line is the same as in Fig. 6. As shown in Fig. 5, the results of the g -matrix folding model (solid lines) are consistent with the data for stable target nuclei. Hence the model prediction, shown by the solid line in Fig. 9, is expected to be reliable also for ^6He particularly at forward angles $\theta_{c.m.} \lesssim 60^\circ$. At backward angles $\theta_{c.m.} \gtrsim 60^\circ$, the solid line is different from the dashed and dotted lines. This means that inclusion of the medium effect and usage of the phenomenological ^4He density seem to be necessary there.




FIG. 7: (Color online) R dependence of microscopic optical potential for $p+{}^6\text{He}$ scattering at $E = 71$ MeV. Solid (dotted) line denotes the potential of the g -matrix folding model with the phenomenological (HF) ${}^4\text{He}$ density. Dash line is the potential of the t -matrix folding model with the phenomenological ${}^4\text{He}$ density.




FIG. 8: (Color online) Potential search for $p+{}^6\text{He}$ scattering at $E = 71$ MeV. The calculations are done with the g -matrix folding model with the phenomenological ${}^4\text{He}$ density. The solid line represents the best-fit result with $(f_V^{\text{CE}}, f_W^{\text{CE}}, f_V^{\text{LS}}, f_W^{\text{LS}}) = (0.7, 0.7, 0.9, 4.0)$, whereas the dashed line corresponds to the original result. Experimental data are taken from Ref. [8, 55].




FIG. 9: (Color online) Angular distributions of the elastic cross section $d\sigma/d\Omega$ (a) and the vector analyzing power A_y (b) for proton scattering from ${}^6\text{He}$ at $E = 200$ MeV. See Fig. 6 for the definition of lines.

C. A dependence of microscopic optical potential

The g -matrix folding model is successful in reproducing the data on polarized proton scattering for targets of $A = 12$ –208. In general, the g -matrix folding model is considered to be less accurate for light targets such as ${}^4, {}^6\text{He}$, since the g -matrix interaction is evaluated in nuclear matter and the local density approximation used seems to be more reliable for heavier targets. Nevertheless, the agreement of the model calculations with the data is still reasonable for light targets ${}^4, {}^6\text{He}$. We then derive global properties of the local microscopic optical potential for proton scattering at 65 MeV. The potential is calculated with the g matrix folding model from the phenomenological target density for $A \leq 40$ and the HF density for $A > 40$.

Since the microscopic optical potential does not have any simple shape, we consider the volume integral J and the RMS radius $\langle R^2 \rangle^{1/2}$ for each part of U :

$$J = \int F(R) d\mathbf{R}, \quad \langle R^2 \rangle^{1/2} = \sqrt{\int R^2 F(R) d\mathbf{R} / J}, \quad (15)$$

where $F(R)$ represents each part of U . As shown in Figs. 3 and 7, W_{LS} is much smaller than V_{LS} and hence yields no significant effect on the present systems. Furthermore, the meaning of the volume integral and the RMS radius is not clear for W_{LS} , since W_{LS} is an oscillating function of R . For these reasons, we do not consider the volume integral and the RMS radius of W_{LS} here.

Figure 10 shows the RMS radius of U as a function of A . For each part of U , A dependence of the RMS radius is fitted for stable targets of $A = 4-208$ by a curve $a_1A + a_2A^{2/3} + a_3A^{1/3} + a_4$; see Table I for the parameter set (a_1, a_2, a_3, a_4) . The fitting line is referred to as the stable-nucleus line. For each panel, the lower figure shows the ratio of each RMS radius to the stable-nucleus line. Since ${}^6\text{He}$ has a halo structure, the ${}^6\text{He}$ RMS radius is enlarged by the structure from the stable-nucleus line by $\sim 18\%$ for V_{CE} , $\sim 25\%$ for W_{CE} and $\sim 12\%$ for V_{LS} . Large enhancement of the ${}^6\text{He}$ RMS radius in W_{CE} means that the reaction cross section is a good tool of searching light halo nuclei and determining their radii.




FIG. 10: (Color online) A dependence of the RMS radius of the microscopic optical potential at 65 MeV. Three panels correspond to V_{CE} , W_{CE} and V_{LS} . See Table I for the parameter set of the stable-nucleus (dotted) line.

	a_1	a_2	a_3	a_4
V_{CE}	-0.0023	0.0104	0.8508	1.0578
W_{CE}	-0.0076	0.0473	0.8732	1.332
V_{LS}	-0.0033	0.0137	1.2305	-0.0682

TABLE I: Parameter fitting for A dependence of the RMS radius of U . All the parameters are presented in units of fm.

We consider the reason why the enhancement of ${}^6\text{He}$ RMS radius is larger in V_{CE} than in V_{LS} . For this purpose, we show A dependence of RMS radii of ρ and $d\rho/dr$ in Fig. 11. In the short-range limit of the g -matrix interaction, the shape of V_{CE} (V_{LS}) is approximately determined by that of ρ ($d\rho/dr$). Enhancement of ${}^6\text{He}$ RMS radius from the stable-nucleus line is smaller for $d\rho/dr$ than for ρ , as shown in Fig. 11; see Table II for the parameter set of the stable-nucleus line (dotted line).




FIG. 11: (Color online) A dependence of the RMS radius of ρ (upper panel) and $d\rho/dr$ (lower panel). See Table II for the parameter set of the stable-nucleus (dotted) line.

Figure 12 shows the volume integral of U as a function of A . The volume integral is also enhanced at $A = 6$ from the stable nucleus line by the halo structure of ${}^6\text{He}$. This enhancement is larger in W_{CE} than in V_{CE} . In V_{LS} , meanwhile, the volume integral is suppressed at $A = 6$ from the stable nucleus line. The halo structure of ${}^6\text{He}$ makes ρ broader and hence suppresses $d\rho/dr$, and consequently weakens the strength of V_{LS} .

	a_1	a_2	a_3	a_4
ρ	-0.0074	0.0378	0.9578	-0.0022
$d\rho/dr$	-0.0079	0.0402	1.2478	-0.2322

TABLE II: Parameter fitting for A dependence of the RMS radii of ρ and $d\rho/dr$. All the parameters are presented in units of fm.

	a_1	a_2	a_3	a_4
V_{CE}	301.28	-59.695	272.22	-125.78
W_{CE}	81.082	92.201	301.22	-439.22
V_{LS}	0.093	6.24	95.645	105.11

TABLE III: Parameter fitting for A dependence of the volume integral of U . All the parameters are presented in units of MeV fm³.

Finally we show A dependence of σ_R in Fig. 13 for proton scattering at $E = 65$ MeV. The reaction cross section is enhanced for ${}^6\text{He}$ by $\sim 35\%$ from the stable-nucleus line (dotted line); see Table IV for the parameter set. This enhancement is a result of the corresponding enhancement in the RMS radius and the volume integral of W_{CE} .

Throughout all the analyses in this subsection, one can find that the broadening of ${}^6\text{He}$ density due to the halo structure (weak-binding nature) yields significant effects on U . The effect is different between the central and spin-orbit forces. For the central force, it enhances the strength and the RMS radius by about 20% $\sim$ 40%. For V_{LS} , meanwhile, the broadening enlarges the RMS radius by about 10% but weakens the strength by about 20%.

D. Isotope dependence of microscopic optical potential

The density broadening due to the weak-binding nature appears not only in ${}^6\text{He}$ but also in the general unstable nuclei where the one-nucleon or the two-nucleon separation energy is small. In addition to the weak-binding nature, the nuclear deformation also makes the diffuseness of density larger, i.e., induces the density broadening.

The success of the local-potential version of the Melbourne g -matrix folding model for ${}^{20-32}\text{Ne} + {}^{12}\text{C}$ scattering and proton scattering from stable nuclei at 65 MeV may indicate that the model is reliable also for $p + {}^{20-32}\text{Ne}$

	a_1	a_2	a_3	a_4
σ_R	-0.6797	45.534	122.34	-170.78

TABLE IV: Parameter fitting for A dependence of σ_R . All the parameters are presented in units of mb.




FIG. 12: (Color online) A dependence of the volume integral of the microscopic optical potential at 65 MeV. See Table III for the parameter set of the stable-nucleus (dotted) line.

scattering at 65 MeV. We then analyze this scattering with the model in order to investigate the weak-binding and the deformation effect on U and σ_R . The analysis is interesting particularly for ${}^{31}\text{Ne}$, since it is a one-neutron halo nucleus with large deformation.

Figure 14 shows isotope dependence of σ_R for $p + {}^{20-32}\text{Ne}$ scattering at 65 MeV. The dotted line denotes the stable-nuclei line for σ_R . The cross symbols represent results of the g -matrix folding model with the spherical HF densities, whereas the closed circles correspond to those with the AMD densities in which Ne isotopes are deformed. The deviation of the cross symbols from the dotted line represents the weak-binding effect, whereas that of the closed circles from the cross symbols shows




FIG. 13: (Color online) A dependence of the reaction cross section for proton scattering at $E = 65$ MeV. See Table IV for the parameter set of the stable-nucleus (dotted) line.

the deformation effect. These effects are more apparent in the lower panel where σ_R for ${}^4\text{Ne}$ is normalized by that for stable nuclei with the same A . At $20 \leq A \leq 23$, the deformation effect is more significant than the weak-binding effect. Since the weak-binding effect becomes large as A increases, the effect is more important than the deformation effect at $24 \leq A \leq 30$. For $A = 31, 32$, Ne isotopes have no bound state in the spherical HF calculation. In this sense, both the deformation and the weak-binding effect are important there.

For ${}^{31}\text{Ne}$, the one-neutron separation energy is quite small. We then do a RGM-AMD calculation to make a tail correction to the AMD density. The closed square denotes a result of the RGM-AMD density, and the deviation of the closed square from the stable-nucleus line represents the net effect of the halo structure and the deformation. This effect is detectable from the measurements of σ_R or the one-neutron removal cross section. Actually, this was done for the ${}^{31}\text{Ne} + {}^{12}\text{C}$ system at 240 MeV/nucleon [24–26, 56, 57].

Figures 15 and 16 show the volume integral and the RMS radius of U for $p + {}^{20-32}\text{Ne}$ scattering at 65 MeV, respectively. The weak-binding nature and the nuclear deformation play the same role in U as in σ_R . Actually, the deformation effect is more important than the weak-binding effect at $20 \leq A \leq 23$, whereas the latter is more significant than the former at $24 \leq A \leq 30$. The density broadening due to the weak-binding nature and the deformation in Ne isotopes yields the same effect on U as that due to the halo structure (weak-binding nature) in ${}^6\text{He}$. For ${}^{31}\text{Ne}$ as a typical case, it suppresses the volume integral by about 10% and enlarges the RMS radius by about 5% for V_{LS} , whereas it enlarges both the volume integral and the RMS radius by about 10% for V_{CE} and W_{CE} . The imaginary central part W_{CE} is more sensitive to the density broadening than the other parts. Thus σ_R is a good quantity to investigate the density broadening.




FIG. 14: (Color online) Isotope dependence of the reaction cross section for $p + {}^{20-32}\text{Ne}$ scattering at 65 MeV. See Table IV for the parameter set of the stable-nucleus (dotted) line.

IV. SUMMARY

We have derived the local microscopic optical potential systematically for polarized proton scattering at 65 MeV, using the local-potential version of the Melbourne g -matrix folding model that was successful in reproducing nucleus-nucleus scattering [24–26]. The local microscopic optical potential for proton scattering well reproduces the experimental data on stable targets systematically and has a geometry similar to the phenomenological optical potentials such the Koning-Delaroche [4] and the Dirac phenomenology [5, 6] one. Also for $p + {}^6\text{He}$ scattering at 71 MeV, the g -matrix folding model yields better agreement than the t -matrix folding model. Therefore we can say that the local-potential version of the Melbourne g -matrix folding model is a reliable model to describe both nucleon-nucleus and nucleus-nucleus scattering.

The local microscopic optical potential U is also quite useful in many applications. Furthermore, the local nature makes it easy to clarify global properties of U over both stable and unstable target nuclei. As target nuclei, we have then considered ${}^6\text{He}$ and neutron-rich Ne isotopes in addition to stable nuclei of mass number $A = 4$ –208. Particularly, it is interesting to compare properties of U in ${}^6\text{He}$ and ${}^{31}\text{Ne}$ with general properties of U in stable nuclei, since ${}^6\text{He}$ is a typical two-neutron halo nucleus and ${}^{31}\text{Ne}$ is a one-neutron halo nucleus with large deformation. Throughout the global analyses, we have clarified A - and isotope dependence of U . The target density is broadened in ${}^6\text{He}$ by the weak-binding nature and in neutron-rich Ne isotopes by both the weak-binding nature and the nuclear deformation. For the central part of U the density broadening enlarges both the strength and the root-mean-square radius, but for the real spin-orbit part of U it enlarges the root-mean-square radius but weakens the strength. The broadening effect is conspicuous for halo nuclei such as ${}^6\text{He}$ and ${}^{31}\text{Ne}$. The statement mentioned above is true also for intermediate energies




FIG. 15: (Color online) A dependence of the volume integral of the microscopic optical potential for $p+^{20-32}\text{Ne}$ scattering at 65 MeV. See Table III for the parameter set of the stable-nucleus (dotted) line.

such as 200 MeV.

Acknowledgments

The authors thank T. Furumoto, T. Matsumoto, M. Kohno, K. Ogata, T. Uesaka and S. Sakaguchi for useful discussions. This work is supported in part by Grant-in-Aid for Scientific Research (No. 244137) from Japan Society for the Promotion of Science (JSPS). The numerical calculations of this work were performed on the computing system in Research Institute for Information




FIG. 16: (Color online) A dependence of the RMS radius of the microscopic optical potential for $p+^{20-32}\text{Ne}$ scattering at 65 MeV. See Table I for the parameter set of the stable-nucleus (dotted) line.

Technology of Kyushu University.

Appendix A: Definitions of quantities

and

The explicit forms of $\delta_0^{(\alpha)}(R, s)$ and $\rho_1^{(\alpha)}(R, s)$ are

$$\begin{aligned} \delta_0^{(\alpha)}(R, s) &= \frac{1}{2} \int_{-1}^{+1} d\omega \frac{g_{\text{LS}, p\alpha}^{\text{EX}}(s; \rho_\alpha)}{x} \\ &\times \left\{ \frac{3}{k_F^{(\alpha)}(x)s} j_1(k_F^{(\alpha)}(x)s) \frac{d}{dx} \rho_\alpha(x) \right|_{x=\sqrt{R^2+s^2/4+Rs\omega}} \\ &+ s \rho_\alpha(x) \frac{d}{dx} k_F^{(\alpha)}(x) \Big|_{x=\sqrt{R^2+s^2/4+Rs\omega}} \\ &\times \frac{d}{dy} \left[\frac{3}{y} j_1(y) \right] \Big|_{y=k_F^{(\alpha)}(x)s} \right\}, \end{aligned} \quad (\text{A1})$$

$$\begin{aligned} \rho_1^{(\alpha)}(R, s) &= \frac{1}{2} \int_{-1}^{+1} d\omega \omega g_{\text{LS}, p\alpha}^{\text{EX}}(s; \rho_\alpha) \frac{3}{k_F^{(\alpha)}(x)s} \\ &\times j_1(k_F^{(\alpha)}(x)s) \rho_\alpha(y) \Big|_{y=\sqrt{R^2+s^2/4+Rs\omega}}, \end{aligned} \quad (\text{A2})$$

where

$$k_F^{(\alpha)} = (3\pi^2 \rho_\alpha)^{1/3}, \quad (\text{A3})$$

$$g_{\text{LS}, pp}^{\text{DR,EX}}(s; \rho_p) = g_{\text{LS}}^{11}, \quad (\text{A4})$$

$$g_{\text{LS}, pn}^{\text{DR,EX}}(s; \rho_n) = \frac{1}{2} (\pm g_{\text{LS}}^{10} + g_{\text{LS}}^{11}). \quad (\text{A5})$$

-
- [1] M. Kamimura, M. Yahiro, Y. Iseri, Y. Sakuragi, H. Kameyama, and M. Kawai, Prog. Theor. Phys. Suppl. **89**, 1 (1986).
 - [2] N. Austern, Y. Iseri, M. Kamimura, M. Kawai, G. Rawitscher, and M. Yahiro, Phys. Rep. **154**, 125 (1987).
 - [3] M. Yahiro, K. Ogata, T. Matsumoto, and K. Minomo, Prog. Theor. Exp. Phys. **2012**, 01A206 (2012).
 - [4] A. J. Koning and J. P. Delaroche, Nucl. Phys. A **713** 231 (2003).
 - [5] S. Hama, B. C. Clark, E. D. Cooper, H. S. Sherif, and R. L. Mercer, Phys. Rev. C **41**, 2737 (1990).
 - [6] E. D. Cooper, S. Hama, B. C. Clark, and R. L. Mercer, Phys. Rev. C **47**, 297 (1993).
 - [7] C. M. Perey and F. G. Perey, At. Data Nucl. Data Tables **17**, 1 (1976).
 - [8] T. Uesaka *et al.*, Phys. Rev. C **82**, 021602(R) (2010); S. Sakaguchi *et al.*, Phys. Rev. C **84**, 024604 (2011).
 - [9] K. Kaki, Y. Suzuki, and R. B. Wiringa, Phys. Rev. C **86**, 044601 (2012).
 - [10] G. Bertsch, J. Borysowicz, H. McManus, and W. G. Love, Nucl. Phys. A **284**, 399 (1977).
 - [11] J. -P. Jeukenne, A. Lejeune, and C. Mahaux, Phys. Rev. C **16**, 80 (1977); J. -P. Jeukenne, A. Lejeune, and C. Mahaux, Phys. Rep. **25**, 83 (1976).
 - [12] F. A. Brieva and J. R. Rook, Nucl. Phys. A **291**, 299 (1977); *ibid.* 291, 317 (1977); *ibid.* 297, 206 (1978).
 - [13] G. R. Satchler and W. G. Love, Phys. Rep. **55**, 183 (1979).
 - [14] G. R. Satchler, "Direct Nuclear Reactions", Oxford University Press, (1983).
 - [15] N. Yamaguchi, S. Nagata, and T. Matsuda, Prog. Theor. Phys. **70**, 459 (1983); N. Yamaguchi, S. Nagata, and J. Michiyama, Prog. Theor. Phys. **76**, 1289 (1986).
 - [16] L. Rikus, K. Nakano, and H. V. Von Geramb, Nucl. Phys. A **414**, 413 (1984); L. Rikus, and H. V. Von Geramb, Nucl. Phys. A **426**, 496 (1984).
 - [17] K. Amos, P. J. Dortmans, H. V. Von Geramb, S. Karataglidis, and J. Raynal, in *Advances in Nuclear Physics*, edited by J. W. Negele and E. Vogt (Plenum, New York, 2000) Vol. 25, p. 275.
 - [18] T. Furumoto, Y. Sakuragi, and Y. Yamamoto, Phys. Rev. C **78**, 044610 (2008); *ibid.*, **79**, 011601(R) (2009); *ibid.*, **80**, 044614 (2009).
 - [19] D. T. Khoa, W. von Oertzen, H. G. Bohlen, and S. Ohkubo, J. Phys. G **34**, R111 (2007).
 - [20] K. Minomo, K. Ogata, M. Kohno, Y. R. Shimizu, and M. Yahiro, J. Phys. G **37**, 085011 (2010) [arXiv:0911.1184 [nucl-th]].
 - [21] K. M. Watson, Phys. Rev. **89**, 575 (1953).
 - [22] A. K. Kerman, H. McManus, and R. M. Thaler, Ann. Phys. **8**, 551 (1959).
 - [23] M. Yahiro, K. Minomo, K. Ogata, and M. Kawai, Prog. Theor. Phys. **120**, 767 (2008).
 - [24] K. Minomo, T. Sumi, M. Kimura, K. Ogata, Y. R. Shimizu, and M. Yahiro, Phys. Rev. C **84**, 034602 (2011).
 - [25] K. Minomo, T. Sumi, M. Kimura, K. Ogata, Y. R. Shimizu, and M. Yahiro, Phys. Rev. Lett. **108**, 052503 (2012).
 - [26] T. Sumi *et al.*, Phys. Rev. C **85**, 064613 (2012).
 - [27] M. Kimura and H. Horiuchi, Prog. Theor. Phys. **111**, 841 (2004).
 - [28] M. Kimura, Phys. Rev. C **75**, 041302(R) (2007).
 - [29] J. F. Berger, M. Girod, and D. Gogny, Comput. Phys. Commun. **63**, 365 (1991).
 - [30] H. de Vries, C. W. de Jager, and C. de Vries, At. Data Nucl. Data Tables **36**, 495 (1987).
 - [31] H. F. Arellano, F. A. Brieva, and W. G. Love, Phys. Rev. C **52**, 301 (1995).
 - [32] R. Machleidt, K. Holinde, and Ch. Elster, Phys. Rep. **149**, 1 (1987).
 - [33] R. P. Singhal, M. W. S. Macauley, and P. K. A. De Witt Huberts, Nucl. Instr. and Meth. **148**, 113 (1978).
 - [34] W. G. Love and M. A. Franey, Phys. Rev. C **24**, 1073 (1981); M. A. Franey and W. G. Love, Phys. Rev. C **31**, 488 (1985).
 - [35] K. Imai *et al.*, Nucl. Phys. A **325**, 397 (1979).
 - [36] K. Fukunaga *et al.*, Nucl. Phys. A **456**, 48 (1986).
 - [37] S. Kato *et al.*, Phys. Rev. C **31**, 1616 (1985).
 - [38] M. Ieiri *et al.*, Nucl. Instr. and Meth. A **257**, 253 (1987).
 - [39] H. Sakaguchi *et al.*, Phys. Lett. B **89**, 40 (1979).
 - [40] H. Ejiri *et al.*, Phys. Rev. C **24**, 2001 (1981).
 - [41] T. Noro *et al.*, Nucl. Phys. A **366**, 189 (1981).

- [42] H. Sakaguchi *et al.*, Phys. Lett. B **99**, 92 (1981).
- [43] A. M. Sourkes, A. Houdayer, W. T. H. van Oers, R. F. Carlson, and R. E. Brown, Phys. Rev. C **13**, 451 (1976).
- [44] A. Ingemarsson *et al.*, Nucl. Phys. A **653**, 341 (1999).
- [45] R. F. Carlson *et al.*, Nucl. Phys. A **445**, 57 (1985).
- [46] N. E. Davison *et al.*, Nucl. Phys. A **290**, 45 (1977).
- [47] J. J. H. Menet, E. E. Gross, J. J. Malanify, and A. Zucker, Phys. Rev. C **4**, 1114 (1971).
- [48] L. G. Votta, P. G. Roos, N. S. Chant, and R. Woody, III, Phys. Rev. C **10**, 520 (1974).
- [49] T. Matsumoto, E. Hiyama, K. Ogata, Y. Iseri, M. Kamimura, S. Chiba, and M. Yahiro, Phys. Rev. C **70**, 061601(R) (2004).
- [50] T. Matsumoto, T. Egami, K. Ogata, Y. Iseri, M. Kamimura, and M. Yahiro, Phys. Rev. C **73**, 051602(R) (2006).
- [51] G. A. Moss *et al.*, Phys. Rev. C **21**, 1932 (1980).
- [52] H. O. Meyer, P. Schwandt, G. L. Moake, and P. P. Singh, Phys. Rev. C **23**, 616 (1981).
- [53] H. Seifert *et al.*, Phys. Rev. C **47**, 1615 (1993).
- [54] L. Lee *et al.*, Phys. Lett. B **205**, 219 (1988).
- [55] A. A. Korshennikov *et al.*, Nucl. Phys. A **617**, 45 (1997).
- [56] T. Nakamura *et al.*, Phys. Rev. Lett. **103**, 262501 (2009).
- [57] M. Takechi *et al.*, Phys. Lett. B **707**, 357 (2012).
- [58] T. Furumoto, W. Horiuchi, M. Takashina, Y. Yamamoto, and Y. Sakuragi, Phys. Rev. C **85**, 044607 (2012).